



Admission Standard Orders Atrial Fibrillation

Admitting Physician: _____ Date: _____ Time: _____

Admit to Inpatient: [] Med/Surg [] Telemetry [] ICU

Place in Observation Services: [] Med/Surg [] Telemetry [] Other _____

Diagnosis: [] Atrial Fibrillation [] Continuous [] Paroxysmal [] Other: _____

Condition: [] Stable [] Guarded [] Critical [] Good [] Fair [] Poor

Consult: [] Cardiology: _____ [] Other: _____

Allergies: [] _____

Code Status: [] Full [] DNR (see DNR order)

Vital Signs: [] Per unit protocol [] Every shift [] every _____ hours

[] Other: _____

Activity: [] Bed rest [] Up in chair [] Bedside commode [] Ambulate ad lib [] Bathroom privileges

Nursing: [] Daily weight [] Intake & Output [] Foley to drainage
[] Pulse oximeter [] every shift [] twice a day [] Continuous overnight monitor
[] Compression Hose (TED)
[] Sequential Compression Device (SCD)
[] Glucose checks AC and at bedtime or every _____ hrs
[] Other: _____

Diet: [] Regular [] NPO [] 2 gm low sodium [] Dietary Consult
[] Clear liquid [] Full liquid [] Cardiac [] Carbohydrate Controlled
[] Other: _____

Fluids: [] Intravenous: _____ @ _____ ml/hr
[] Saline lock
[] Other: _____

Oxygen: [] Nasal Cannula _____ L/min, adjust to keep O2 sat greater than 92%. May DC if O2 sat greater than 93% on RA.
[] Venti Mask _____ % FIO2 [] 100% NRB
[] Other: _____

Protocols (if available): All protocol orders must be in chart

- [] Weight Based Heparin Dosing Protocol (when patient placed on protocol, all associated labs and monitoring is included)
- [] Insulin Infusion Protocol (ICU patients only)

Labs: [] Troponin (STAT repeat 90 minutes, and 12 hours)
[] BMP [] CMP [] BNP [] CPK [] Myoglobin [] Magnesium [] ABG
[] Fasting lipid profile [] TSH [] Free T4 [] UA [] CBC
[] Digoxin level
[] Other Labs: _____

Studies: (If LVEF less than 40%, implement CHF Protocol)

- [] C XR: [] Portable [] PA/Lateral
- [] EKG
- [] STAT EKG PRN with chest pain or palpitations
- [] Echocardiogram Dr. _____ to read
- [] 24 hour Holter Monitor
- [] Other: _____

NURSE NOTED	DATE	TIME	DATE / TIME	PHYSICIAN SIGNATURE OR AUTHENTICATION
24 HR. CHART CHECK BY NURSE	DATE	TIME		



Account Number:	MR Number:
Patient Name:	
Admit Date:	

Davis Hospital AND MEDICAL CENTER 1600 West Antelope Drive - Layton, UT 84041 (801) 807-1000	DOB	Age	Sex	HT	WT	RM-BD	PT	SVC	FC
	Allergies:								
	Attending Physician Name:								

Medications: (Check the appropriate box)

- Diltiazem (Cardizem) 0.25 mg/kg loading dose (max 0.20 mg) IV over 5 minutes, then start drip at 10 mg/hr
If patient older than 65 years or actual body weight less than 60 kg., reduce loading dose to 15 mg and start drip at 5 mg/hr.
- Diltiazem CD (Cardizem CD) 120 mg 180 mg 240 mg 300 mg PO daily
- Metoprolol (Lopressor) 12.5 mg PO twice daily 25 mg PO twice daily 50 mg PO twice daily 100 mg PO twice daily
- Metoprolol XL (Toprol XL) 25 mg PO daily 50 mg PO daily 100 mg PO daily 200 mg PO daily
- Digoxin (Lanoxin) 0.5 mg IV bolus, followed by 0.25 mg in 6 hrs and 12 hrs. Then 0.125 mg daily 0.25 mg PO daily.
- Amiodarone (Cordarone) 150 mg in 100ml Dextrose 5% Water IV, infuse over 10 minutes, then 1 mg per minute IV for 6 hours then 0.5 mg per minute IV for 18 hrs.
- Propafenone (Rythmol) 150 mg PO 225 mg PO 300 mg PO, every 8 hours
- Procainamide (Procan, Procanbid) 15 mg/kg IV loading dose at 20 mg per minutes, then 2 mg 3 mg 4 mg per minute.
- Esmolol (Brevibloc) 500 micrograms/kg bolus infused over 1 minute, followed by 50 micrograms/kg per minute maintenance infusion
- Enoxaparin (Lovenox) 40 mg subcutaneous every 24 hrs (pharmacy to adjust for renal dysfunction)
- Anticoagulant protocol per pharmacy.
- Percocet 5/325 mg PO every 4 hrs PRN for moderate pain
- Morphine Sulfate 2 mg IV every 4 hrs PRN severe pain (unless patient is allergic to codeine or morphine)
- Lorazepam (Ativan) 0.5 mg PO every 6 hrs PRN 1 mg PO PRN every 6 hrs PRN as needed for mild to moderate anxiety
- Famotidine (Pepcid) 20 mg IV twice daily if NPO, otherwise 20 mg PO twice daily
- Pantoprazole (Protonix) 40 mg IV once daily in NPO, otherwise Prilosec 20 mg PO daily
- Nicotine _____ mg topical patch apply daily
- Temazepam (Restoril) 15 mg PO at bedtime PRN sleep. May repeat does once in one hour if no results
- Ondansetron (Zofran) 4 mg IV every 8 hrs PRN for nausea vomiting
- Percocet 5/325 mg PO every 4 hrs PRN for moderate pain



Standard Medication:

- Acetaminophen (Tylenol) 650 mg PO every 4 hr PRN mild pain (not to exceed 4 grams per 24 hours)
- Acetaminophen (Tylenol) 650 mg PO every 4 hr PRN temp more than 101 F (not to exceed 4 grams per 24 hours)
- Docusate sodium (Colace) 100 mg PO qHS
- MOM 30 ml PO every HS PRN constipation
- Maalox 30 ml PO every 4 hrs PRN heartburn

Additional Meds:

- _____
- _____
- _____
- _____
- _____
- _____
- _____

NURSE NOTED	DATE	TIME	DATE / TIME	PHYSICIAN SIGNATURE OR AUTHENTICATION
24 HR. CHART CHECK BY NURSE	DATE	TIME		

 	Account Number:	MR Number:							
Patient Name:									
Admit Date:									
Davis Hospital AND MEDICAL CENTER 1600 West Antelope Drive - Layton, UT 84041 (801) 807-1000	DOB	Age	Sex	HT	WT	RM-BD	PT	SVC	FC
Allergies:									
Attending Physician Name:									